LADY ROSETTA

Cultivation advice

Purpose: Crisp

Characteristics:

Maturity	7
Colour of skin	R
Color of flesh	6
Shape of tuber	R
Depth of the eyes	6
Regularity shape of tuber	8.5
Size of tubers	7
Grading	8.5
Dry matter percentage	25
Suitability for cooking	7C
Dormancy	5.5

Comments:
9 = positive .... 1 = negative
Yellowness of flesh: 8 = yellow . . . . 4 = white
Colour of skin: L=light, D=dark, W=white, G=yellow, R=red
Shape of tuber: R = Round O = Oval L = Long
Cooking type: A = Firm . . . . D = Very mealy

Planting distance:

Size	Rowspace 75 cm	Plants/hectare (x 1000)
25/28	16	83,3
28/35	19	70,2
35/50	27	49,4
35/55	29	46,0
50/55	33	40,4
50/60	35	38,1
55/60	40	33,3

Fertilisation (kg/ha):

depending on the soil reserve



				Total
Nitrogen:	(N)	200	75	275
Phosphate:	(P_2O_5)	110		110
Potassium:	(K_2O)	230	100	330

Storage:



8 °C
4 months

Wait for eyes waking up before planting
Start late blight control in time
Wait for a set skin before harvest
After harvest, dry the potatoes by ventilation
Avoid black spot

Resistance:

Leaf roll	6
A-virus	R
X-virus	R
Y-virus	7.5
Yntn-virus	8
Foliage blight	3
Tuber blight	5.5
Wart disease	wv1
Potato cyst nematode	Ro1,4
Common scab	6
Spraing	7.5
Bruising	5
Secondary growth	7.5

Comments:
R = resistant .... 1 = very susceptible
Ro(x) = Globodera Rostochiensis pathotype (x)
Pa(x) = Globodera Pallida pathotype (x)
(wv)O1 = (little susceptible) wart disease fysio 1

C. Meijer B.V. will not be responsible for possible negative effects originating from this advice.